

2011. Thanks also go to all the 2013 autumn season volunteers, particularly Sebastian Seibold, Viktoria Mader and Christine Thorn, and to Nial Moores, Kiyooki Ozaki and Yoshiki Watabe for helpful comments. This study was supported by the Oriental Bird Club, Deutsche Ornithologen-Gesellschaft e.V. and Förderkreis für Allgemeine Naturkunde (Biologie) e.V.

References

- BirdLife International (2015) Species factsheet: *Parus venustus*. Downloaded from <http://www.birdlife.org> on 10/08/2015.
- Choi C. Y., Park J. G., Moores, N., Kim E. M., Kang C. W., Nam H. Y. & Kim S. M. (2011) The recent increase in the Redbilled Starling *Sturnus sericeus* in the Republic of Korea. *Forktail* 27: 89–91.
- Dai C., Chen K., Zhang R., Yang X., Yin Z., Tian H., Zhang Z., Hu Y. & Lei F. (2010) Molecular phylogenetic analysis among species of Paridae, Remizidae and Aegithalidae based on mtDNA sequences of COI and cyt b. *Chinese Birds* 1: 112–123.
- Gosler, A. & Clement, P. (2016). Yellow-bellied Tit (*Pardaliparus venustus*). In J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie & E. de Juana, eds. *Handbook of the birds of the world alive*. Barcelona: Lynx Edicions. (retrieved from <http://www.hbw.com/node/59875> on 07/08/2016).
- Harrap, S. & Quinn, D. (1996) *Tits, nuthatches & treecreepers*. London: Christopher Helm.
- Hebert, P. D. N., Stoeckle, M. Y., Zemlak, T. S. & Francis, C. M. (2004) Identification of birds through DNA barcodes. *PLoS Biol.* 2: e312.
- Heim, W. (2016) A survey of breeding waterbird communities on lakes and other waterbodies on the middle reaches of the Amur River valley near Blagoveshensk, Amur province, Far East Russia. *BirdingASIA* 25: 98–103.
- Heim, W. & Smirenski, S. M. (2013) The Amur bird project at Muraviovka Park in Far East Russia. *BirdingASIA* 19: 31–33.
- Ikenaga, H., Kawakami, K. & Yanagisawa, N. (2014) A note on the newly accepted species and subspecies in the check-list of Japanese birds, seventh revised edition. *Japanese J. Orn.* 63: 96–149.
- Martens, J., Tietze, D. T. & Sun Y. H. (2006) Molecular phylogeny of *Parus* (*Periparus*), a Eurasian radiation of tits (Aves: Passeriformes: Paridae). *Zool. Abhandl. Staatl. Mus. Tierk. Dresden* 55: 103–120.
- Moores, N. (2005) Yellow-bellied Tit *Parus venustus*. Downloaded from http://www.birdskorea.org/Birds/Significant_Records/Rarity_Reports/BK-RR-Yellow-bellied-Tit-2005-10.shtml. on 07/08/2016.
- Moores, N., Kim A. & Kim R. (2014) *Status of birds 2014*. Available at <http://www.birdskorea.org/Habitats/Yellow-Sea/YSBR/Downloads/Birds-Korea-Status-of-Birds-2014.pdf>.
- Tamura, K., Stecher, G., Peterson, D., Filipowski, A. & Kumar, S. (2013) MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol. Biol. Evol.* 30: 2725–2729.
- Williams, M. D., Carey, G. J., Duff, D. G. & Weishu X. (1992) Autumn bird migration at Beidaihe, China, 1986–1990. *Forktail* 7: 3–55.
- P. FETTING**, Goethestraße 1, 17489 Greifswald, Germany.
- S. THORN**, Field Station Fabrikschleichach, Department of Animal Ecology and Tropical Biology, Biocenter University of Würzburg, Glashüttenstr, 5, 96181 Rauhenebrach, Germany.
- M. PÄCKERT**, Senckenberg Naturhistorische Sammlungen, Königsbrücker Landstraße 159, 01109 Dresden, Germany.
- W. HEIM**, Amur Bird Project, Roseggerstraße 14, 14471 Potsdam AND Animal Ecology, University of Potsdam, Maulbeerallee 1, 11469 Potsdam, Germany. Email: amurbirding@gmx.de (corresponding author).

First breeding record of Slaty Bunting *Emberiza siemsseni*

YUNBIAO HU & YUEHUA SUN

The Slaty Bunting *Emberiza siemsseni* is a range-restricted Chinese endemic, breeding in the highlands of south Gansu province, south Shaanxi province and west and north-east Sichuan province (Zheng 2011). Although not rare and designated as Least Concern by BirdLife International (2015), very little is known about its breeding ecology (Madge 2015). Here, we describe for the first time the Slaty Bunting's eggs, nest, nest-site and breeding behaviour.

On 3 July 2014 we found a Slaty Bunting nest in Lianhuashan Nature Reserve, Kangle county, Gansu province (34.685°N 103.523°E). The nest was located 0.3 m from a small path in shrub-land and was positioned 0.4 m above the ground on a 0.7 m tall spruce sapling. About 50% of the nest site area (defined as a 5 m radius around the nest) was covered by shrubs, dominated by willows *Salix* sp., with an average height of 3.5 m.

When found, the nest contained four eggs. No further eggs were laid, hence incubation was already underway. The eggs were oval in shape and creamy white in colour, with irregular markings slightly concentrated at the large end (Plate 1). We weighed the eggs to the nearest 0.01 g, using a portable digital balance, and measured the length and breadth to the nearest 0.01 mm, using digital vernier calipers. Mean egg size was 17.93 ± 0.38 mm \times 14.10 ± 0.11 mm, with a mean weight of 1.89 ± 0.07 g.

The outer layer of the circular, cup-shaped nest was made mainly of broad leaves and grass-stems, with a lining of soft, thin grass-stems and livestock hairs. The inner diameter of the nest was 57.7 mm and the outer diameter 84.4 mm, with an inner depth of 44.7 mm and total height of 65.5 mm.

We placed an infrared-triggered camera near the nest to monitor parental activities. However, because of rainy weather and the

limitations of the camera, we obtained detailed incubation data only on 10 July. From 06h00 to 18h00 we observed 12 change-overs, and both sexes spent similar times incubating: 349 minutes (male) and 352 minutes (female). Three eggs hatched on 13 July, when the first images were captured of the adults feeding the nestlings, giving an incubation period of at least 10 days. One egg failed to hatch, and it was examined after the nestlings had left the nest: the yoke was still complete and no embryo development was evident, so we deduced that the egg was unfertilised.

Both sexes reared the nestlings, but detailed provisioning rates could not be obtained because of the two-minute intervals between consecutive triggered events on the camera. We watched 10 feeding sessions and observed that the parents mainly fed insects and insect larvae to their nestlings, with some spiders and other small arthropods. Ten days after hatching, the plumage of the nestlings resembled that of the female in colour and appearance (Plate 2). The young birds remained in the same vicinity at least until 29 July because the adults were observed carrying food near the nest site every day.

Recent phylogenetic analyses indicate that the Slaty Bunting is a sister species to Yellow-throated Bunting *E. elegans* (Alström *et al.* 2008). Compared with the Yellow-throated Bunting (Chen *et al.* 2015), our results suggest that the Slaty Bunting has smaller eggs and males may take a greater part in incubation. However, as we only observed one nest of the Slaty Bunting, we must be cautious in any general statement regarding its breeding biology.

Although Slaty Bunting is not a threatened species, its breeding range is relatively narrow; the Lianhuashan Nature Reserve is about 150 km north of the previously known breeding range of



YUNBIAO HU

Plate 1. Nest and eggs of the Slaty Bunting *Emberiza siemsseni*, Lianhuashan Nature Reserve, Kangle county, Gansu province, China, 3 July 2014.



YUNBIAO HU

Plate 2. Slaty Bunting nestlings at 10 days old, Lianhuashan Nature Reserve, 22 July 2014.

the species. By checking the bird species list of the reserve (Sun *et al.* 2008) and many birdwatching notes, we confirmed that it was a new record for Lianhuashan. Between late April and early June 2014, we trapped four male Slaty Buntings; only one male was seen in 2015, on 15 April; in 2016 we trapped one male, on 19 May. We tried to locate more birds by using song playback and mist-nesting, but none were found in either the 2015 or 2016 breeding seasons. These records suggest that the Slaty Bunting may have extended its breeding range to the north in recent years, although more field evidence is needed to verify this. We hope that our findings will stimulate other ornithologists to search for nests and collect more information about the species's breeding biology in its normal breeding range.

Acknowledgements

We thank Chen Lijun for providing details of the 2015 record, and Lou Yingqiang and the staff of Lianhuashan Nature Reserve for their help in the field. This work was funded by National Natural Science Foundation of China (No. 31270468).

References

Alström, P., Olsson, U., Lei F. M., Wang H. T., Gao W. & Sundberg, P. (2008) Phylogeny and classification of the Old World Emberizini (Aves, Passeriformes). *Mol. Phylogenet. Evol.* 47: 960-973.

BirdLife International (2015) Species factsheet: *Emberiza siemsseni*. Downloaded from <http://www.birdlife.org> on 13/12/2015.

Chen W. K., Xie H., Shi X. X. & Fan Q. F. (2015) Breeding notes of Yellow-throated Bunting (*Emberiza elegans*). *Chinese J. Zool.* 50: 621-627. (In Chinese.)

Madge, S. (2015) Slaty Bunting (*Emberiza siemsseni*). *HBW Alive* (retrieved from <http://www.hbw.com/node/61860> on 13/12/2015).

Sun Y. H., Fang Y., Klaus, S., Martens, J., Scherzinger, W. & Swenson, J. E. (2008) *Nature of the Lianhuashan Natural Reserve*. Shenyang: Liaoning Science and Technology Publishing House.

Zheng G. M. (2011) *A checklist on the classification and distribution of the birds of China*. Second Edition. Beijing: Science Press.

Yunbiao HU, Key Laboratory of Animal Ecology and Conservation Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China Email: huyb@ioz.ac.cn

Yuehua SUN, Key Laboratory of Animal Ecology and Conservation Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China Email: sunyh@ioz.ac.cn

Further evidence that wasps prey on nestlings

YI-QIANG FU, MING XIANG, CHI-PING KONG & YONG-HENG WU

Social wasps of the family Vespidae forage for water, pulp, carbohydrates and animal protein. When hunting, they are opportunistic generalists and use a variety of mechanisms to locate and choose prey which they consume directly (Edwards 1980, Richter 2000). There are complicated ecological associations between wasps and birds (Almeida & Anjos-Silva 2015), such as competition for food (Beggs 2001). However, increasing evidence shows that birds are one of the important predators of wasps (Windsor 1976, Henriques & Palma 1998, Almeida & Anjos-Silva 2015), but wasps are also thought to kill nestlings (Moller 1990). Here, we summarise a striking example of wasp–bird interactions.

During the 2015 breeding season, we studied the feeding habits of Red-billed Leiothrix *Leiothrix lutea* at the Laojunshan National Nature Reserve, Sichuan province, south-west China

(28.660–28.727°N 103.960–104.070°E). The reserve lies between 900 and 2,000 m. The climate is temperate (annual average temperature 12.0–14.7°C) with high precipitation (more than 1,500 mm per year) and the characteristic vegetation is evergreen broadleaf forest (Fu *et al.* 2011). Social wasps are common in the lower parts of the reserve.

On the afternoon of 11 August 2015, we recorded the predation of Red-billed Leiothrix nestlings in nest 30-2015, by social wasps. The four nestlings were about three days old and the main sequence of events was recorded using a Canon SX50 HS digital camera as detailed here. At 16h26:31 an adult leiothrix arrived and cleaned the nest. The nestlings were active at the time; the adult left at 16h27:04. Then at 16h32:58, a single wasp arrived, entered the nest and killed the four nestlings; it left at 16h39:05, carrying nestling flesh (Plate 1). At 16h50:21 an adult leiothrix arrived with insects to feed the