

## Phylogeny and Biogeography of *Cedrus* (Pinaceae) Inferred from Sequences of Seven Paternal Chloroplast and Maternal Mitochondrial DNA Regions

CAI-YUAN QIAO<sup>1,2,†</sup>, JIN-HUA RAN<sup>1,†</sup>, YAN LI<sup>1,2</sup> and XIAO-QUAN WANG<sup>1,\*</sup>

<sup>1</sup>State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, The Chinese Academy of Sciences, Beijing 100093, China and <sup>2</sup>Graduate School of the Chinese Academy of Sciences, Beijing 100039, China

Received: 24 December 2006 Returned for revision: 16 March 2007 Accepted: 23 May 2007 Published electronically: 4 July 2007

- **Background and Aims** *Cedrus* (true cedars) is a very important horticultural plant group. It has a disjunct distribution in the Mediterranean region and western Himalaya. Its evolution and biogeography are of great interest to botanists. This study aims to investigate the phylogeny and biogeography of *Cedrus* based on sequence analyses of seven cytoplasmic DNA fragments.
- **Methods** The methods used were PCR amplification and sequencing of seven paternal cpDNA and maternal mtDNA fragments, parsimony and maximum likelihood analyses of the DNA dataset, and molecular clock estimate of divergence times of *Cedrus* species.
- **Key Results** Phylogenies of *Cedrus* constructed from cpDNA, mtDNA and the combined cp- and mt-DNA dataset are identical in topology. It was found that the Himalayan cedar *C. deodara* diverged first, and then the North African species *C. atlantica* separated from the common ancestor of *C. libani* and *C. brevifolia*, two species from the eastern Mediterranean area. Molecular clock estimates suggest that the divergence between *C. atlantica* and the eastern Mediterranean clade at  $23.49 \pm 3.55$  to  $18.81 \pm 1.25$  Myr and the split between *C. libani* and *C. brevifolia* at  $7.83 \pm 2.79$  to  $6.56 \pm 1.20$  Myr.
- **Conclusions** The results, combined with palaeogeographical and palaeoecological information, indicate that *Cedrus* could have an origin in the high latitude area of Eurasia, and its present distribution might result from vicariance of southerly migrated populations during climatic oscillations in the Tertiary and further fragmentation and dispersal of these populations. It is very likely that *Cedrus* migrated into North Africa in the very late Tertiary, while its arrival in the Himalayas would not have been before the Miocene, after which the phased or fast uplift of the Tibetan plateau happened.

**Key words:** *Cedrus*, molecular phylogeny, biogeography, Mediterranean, Himalayas, molecular clock.

### INTRODUCTION

Mountain regions between the fortieth parallels of latitude are of interest to biogeographers due to their importance in the survival of many plant species during the Ice Ages (Hewitt, 2004). The Mediterranean is one of 18 world hot-spots of biodiversity, where approx. 25 000 plant species, of which 50% are endemics, are harboured (Cowling *et al.*, 1996; Myers *et al.*, 2000; Scarascia-Mugnozza *et al.*, 2000; Comes, 2004). Phylogeographical studies have provided strong evidence that some refugia were available in montane Mediterranean during the Quaternary, and also shaped the distribution of modern biota (Svenning, 2003; Comes, 2004; Hellwig, 2004). Unfortunately, the origin of the Mediterranean flora is still obscure (Valcárcel *et al.*, 2003). In Asia, the uplift of the Himalayas and the onset of the Asian monsoon changed the world climate (Griffin, 2002; Liu and Yin, 2002), and made the Tibetan plateau, especially its south-eastern part, an important centre for not only the survival of many arctic plants, when they migrated southwards, but also the speciation of various organisms (Axelrod *et al.*, 1996; Myers *et al.*, 2000). It is of great interest to investigate the floristic relationship between the Mediterranean and the Tibetan plateau.

Moreover, the significant geological and climatic changes in the two regions during the Cenozoic (Mai, 1989; LePage, 2003; Hellwig, 2004) produced many disjunct distributions (Sanmartín, 2003; Comes, 2004; Hellwig, 2004), and thus give us opportunities to test the dispersal and vicariance theories of biogeography.

*Cedrus* (true cedar) is one of 11 commonly accepted genera in Pinaceae, first described by Trew in 1757 (Farjón, 2001). It comprises four species with a highly disjunct distribution in circum-Mediterranean and western Himalayas (Farjón, 1990, 2001), i.e. *Cedrus deodara* (Roxb.) G. Don in the Hindu Kush, Karakoram and Indian Himalayas, *Cedrus libani* A. Rich. in Turkey, Lebanon and Syria, *Cedrus brevifolia* (Hook. f.) Henry in Cyprus, and *Cedrus atlantica* (Endl.) Manetti ex Carrière in North Africa (Algeria, Morocco). The true cedars occurring around the Mediterranean basin were once treated as one species *C. libani*, including four subspecies, namely ssp. *atlantica*, ssp. *brevifolia*, ssp. *stenocoma* in Turkey and ssp. *libani* in Lebanon and Syria (Davis, 1965). Some authors divided *Cedrus* into three species (cf. Scaltsoyiannes, 1998), regarding *C. brevifolia* as a subspecies of *C. libani* (Meikle, 1977). The great debate in species delimitation and phylogeny of *Cedrus* remains unresolved, although previous cytological (Li and Fu, 1995; Bou Dagher-Kharrat *et al.*, 2001), biochemical (Scaltsoyiannes, 1998) and gene

<sup>†</sup> These authors contributed equally to the work.

\* For correspondence. E-mail xiaoq\_wang@ibcas.ac.cn

flow analyses (Fady *et al.*, 2003) contributed greatly to the interpretation of interspecific relationships of true cedars.

In recent years, sequence analysis of multiple genes from different genomes has been successfully used in reconstructing complex phylogenies, meanwhile providing important information to the inference of biogeographical histories of many plant groups (e.g. Baum *et al.*, 1998; Kusumi *et al.*, 2002; Schneeweiss *et al.*, 2004; Winkworth and Donoghue, 2005; Xiang *et al.*, 2005). The combined gene analysis has great advantages in some conifer groups such as Pinaceae (Wang *et al.*, 2000), where chloroplast and mitochondria are paternally and maternally inherited, respectively (Hipkins *et al.*, 1994; Birky, 1995; Mogensen, 1996; Ahuja, 2001). The joint cp- and mt-DNA analysis can detect interspecific gene flow and reticulate evolution. However, no DNA sequence-based phylogeny has been constructed for *Cedrus*, a very important horticultural plant group. On the other hand, the systematic position of *Cedrus* in Pinaceae is also quite controversial, like the difficulty in species delimitation of the genus. Based on leaf arrangement, the presence of short shoots, and the petiole and pulvinus form, *Cedrus* was placed in the subfamily Laricoideae, with *Larix* and *Pseudolarix* (Cheng and Fu, 1978; Krüssmann, 1985). This classification was not supported by the subsequent anatomical (Hart, 1987; Frankis, 1988; Wu and Hu, 1997), immunological (Price *et al.*, 1987) and cytological studies (Li, 1995). In particular, *Cedrus* does not have the same position in molecular phylogenies constructed by different markers such as PCR-RFLP (Tsumura *et al.*, 1995) and DNA sequence (Quinn *et al.*, 2002), and may represent a very distinctive lineage (Wang *et al.*, 2000). The reconstruction of a robust phylogeny of *Cedrus* comprising all species could also shed some light on the origin of the genus.

To resolve interspecific relationships of plants, sequences of intergenic spacers and gene introns are most widely used because of their fast evolutionary rates (Wang *et al.*, 1999; Wei and Wang, 2003; Shaw *et al.*, 2005). Here, the phylogeny of true cedars based on sequences of seven cytoplasmic DNA fragments, including five from the paternal chloroplast genome and two from the maternal mitochondrial genome, is reconstructed, and the intrageneric classification, divergence times and biogeography of *Cedrus* are discussed.

## MATERIALS AND METHODS

### Plant materials

All four species of *Cedrus* were sampled, and each species was represented by two individuals. To investigate whether the molecular markers we used have infraspecific variation, six individuals of *C. libani* that are at least 100 m apart were further analysed. *Pinus armandii* Franchet and *Cathaya argyrophylla* Chun et Kuang, each represented by one individual, were chosen as outgroups based on molecular phylogenies of Pinaceae (Wang *et al.*, 1998, 2000; Quinn *et al.*, 2002). The origins of materials are shown in Table 1.

### DNA extraction, PCR amplification and sequencing

Total DNA was extracted from silica gel-dried needles using the CTAB method following the protocol of Rogers and Bendich (1988) and used as the template in the polymerase chain reaction. The seven cytoplasmic DNA fragments used were chloroplast *trnT-trnF* (Wei and Wang, 2003), *trnC-trnD* (Lee and Wen, 2004), *5'rps12-rpL20*, *psbB-psbH* and *rpL16* intron (Shaw *et al.*, 2005), and mitochondrial *nad1* intron 2 (Song *et al.*, 2002; Won and Renner, 2003) and *nad5* intron 1 (Jaramillo-Correa *et al.*, 2003). These gene regions were amplified using the primers listed in Table 2. The PCR reaction was carried out in a volume of 25 µL containing 5–50 ng of DNA template, 6.25 pmol of each primer, 0.2 mM of each dNTP, 2 mM MgCl<sub>2</sub>, and 0.75 U of *Taq* DNA polymerase. Amplification was conducted in a Tpersonal Thermocycle and T1 Thermocycle (Biometra, Goettingen, Germany). PCR cycles were as follows: one cycle of 4 min at 70 °C, four cycles of 40 s at 94 °C, 20 s at 52–58 °C, and 2–2.5 min at 72 °C, followed by 36 cycles of 20 s at 94 °C, 20 s at 50–56 °C, and 2–2.5 min at 72 °C, with a final extension step for 10 min at 72 °C. PCR products were separated by 1.5 % agarose gel electrophoresis. The band with the right size was cut out and purified using the GFX PCR DNA and Gel Band Purification Kit (Amersham Biosciences, Amersham, UK). Sequencing reactions were performed with the two PCR primers listed above and several internal primers using the DYEnamic Energy Transfer (ET) Terminator Reagent Premix Kit (Amersham Biosciences). The internal primers included c (Wei and Wang, 2003) for the *trnT-trnF* region, *petN2G* (5'-CTTGGGCTGCTTTA ATGGT AG-3', forward), *psbM2GF* (5'-GTAGAGCAGC AATAAATGCAAG-3', forward), *psbM2GR* (5'-CTTGCA TTTATTGCTGCTCTAC-3', reverse) and *petN3G* (5'-ATGGTACGAGGTCCTTCATCC-3', forward) for the *trnC-trnD* region, and *nad5-IF* (5'-GGCTTTAGGGGGC CTTATG-3', forward) for the *nad5* intron 1. *petN2G*, *psbM2GF* and *psbM2GR* were modified from *petN2*, *psbM2* and *psbM2R* of Lee and Wen (2004), respectively. After precipitation in 95 % EtOH and 3 M NaAc (pH 5.2), the sequencing products were separated on a MegaBACE 1000 automatic sequencer (Amersham Biosciences).

### Data analysis

Sequence alignments were made with CLUSTAL X (Thompson *et al.*, 1997) and refined manually for the maximization of sequence homology using BioEdit 5.0.9 (Hall, 1999). Maximum parsimony analysis was performed using PAUP version 4.0b10 (Swofford, 2002) with *Pinus armandii* and *Cathaya argyrophylla* as outgroups (Wang *et al.*, 2000). Branch-and-bound searches were conducted with the MULTREES option. Indels in the alignment induced by the ingroups were coded as 1/0 binary characters, and gaps of different lengths were all treated as single events. All character states were specified as unordered and equally weighted. To evaluate relative robustness of the clades found in the most-parsimonious trees, the non-parametric bootstrap analysis (Felsenstein, 1985) was

TABLE 1. Sources of material

Taxa	Origins	Individuals/vouchers
Ingroups		
<i>Cedrus atlantica</i> (Endl.) Manetti ex Carrière	Natural History Museum, Budapest, Hungary (cultivated)	2/Folly acb 1905, Debreczy-ca1
<i>Cedrus brevifolia</i> (Hook. f.) Henry	Natural History Museum, Budapest, Hungary (cultivated)	1/Debreczy-cb1
	Cyprus	1/Debreczy-cb2
<i>Cedrus deodara</i> (Roxb.) G. Don	Botanic Garden, Institute of Botany, Beijing, China (cultivated)	2/Wang-cd1, Wang-cd2
<i>Cedrus libani</i> A. Rich.	Elmali, Antalya, Turkey	7/Wang-cl-15
	Taurus Mt, Turkey	1/Debreczy-cl1
Outgroups		
<i>Pinus armandii</i> Franchet	Botanic Garden, Institute of Botany, Beijing, China (cultivated)	1/Wang 6102
<i>Cathaya argyrophylla</i> Chun et Kuang	Jinfo Mountain, Nanchuan, Sichuan, China	1/WangXQ94513

DNA sequences of these materials determined in the present study have been deposited in GenBank under accession numbers DQ983599–DQ983604 (*nad1*), DQ983605–DQ983610 (*nad5*), DQ983611–DQ983616 (*psbB-psbH*), DQ983617–DQ983622 (*rpL16*), DQ983623–DQ983628 (*5'rpS12-rpL20*), DQ983629–DQ983634 (*trnC-trnD*) and DQ983635–DQ983640 (*trnT-trnF*). All vouchers are deposited in PE (Herbarium, Institute of Botany, the Chinese Academy of Sciences, Beijing).

performed with 1000 replicates using the same search settings. The incongruence length difference test (Farris *et al.*, 1995) was conducted, using the partition homogeneity test in PAUP ver. 4.0b10 (Swofford, 2002), to examine the congruence between different datasets. Test settings were 100 random stepwise additions and 1000 replicates of heuristic search with TBR branch swapping using two outgroups.

Owing to the large length variation in the mitochondrial *nad1* intron 2, only the combined chloroplast dataset was used to estimate divergence times of *Cedrus* species. To estimate branch lengths, sequences of the ingroups were fitted using maximum likelihood (ML) models estimated on the topology of the parsimony trees. Optimal ML models for the combined data were selected using Modeltest 3.06 (Posada and Crandall, 1998). To estimate ages of nodes, first the hypothesis of a molecular clock was tested for the dataset using a Likelihood Ratio Test (LRT) that compared likelihood scores of the ML phylogenetic hypothesis with (L1) and without (L0) the molecular clock enforced. Significance was assessed by comparing two times the difference in log likelihoods to a chi-square distribution with  $n - 2$  degrees of freedom, where  $n$  was the number of ingroups. When LRT showed that the data did not adhere to a molecular clock, non-parametric rate

smoothing (NPRS) (Sanderson, 1997) in the program TreeEdit (<http://evolve.zoo.ox.ac.uk/software/TreeEdit/>; Rambaut and Charleston, 2002) transformed ML trees into ultrametric topologies. All node heights, sum of the branch lengths from a node to a tip, were transformed from units of molecular change into units of time by calibration to a node of known age. In addition, standard errors for estimates of node age were calculated for two nodes by using the non-parametric bootstrap resampling method (100 replicates) in PAUP\*.

Since the NPRS method does not use rate smoothing, penalized likelihood (PL) analysis (Sanderson, 2002) was used additionally to estimate the divergence times of *Cedrus*. The ages of the nodes in the tree were estimated using penalized likelihood rate smoothing under TN algorithm with the program r8s version 1.7 (Sanderson, 2003, available at <http://ginger.ucdavis.edu/r8s>). A cross-validation analysis was performed to obtain the most likely smoothing parameter. Standard errors of the divergence times were estimated by a three-step nonparametric bootstrap procedure (Efron and Tibshirani, 1993): (1) 100 data sets were simulated by using the SEQBOOT program in PHYLIP Version 3.6a2 (Felsenstein, 1993); (2) the matrices were imported into PAUP version 4.0b10 and ML trees were generated; (3) the tree file was processed

TABLE 2. Sequences and references for chloroplast and mitochondrial DNA primers used in the present study

Region	Forward primer	Sequence (5'–3')	Reverse primer	Sequence (5'–3')	Reference
cpDNA					
<i>psbB-psbH</i>	<i>psbB</i> (F)	TCCAAAACTGGGAGATCCAAC	<i>psbH</i> (R)	TCAATGGTCTGTGTAGCCAT	Shaw <i>et al.</i> , 2005*
<i>rpL16</i>	<i>rpL16</i> F	GCTATGCTTAGTGTGCGACTCGTTG	<i>rpL16</i> R	CCYTTCATTCTTCCCCTATGTTG	Small <i>et al.</i> , 1998*
<i>5'rpS12-rpL20</i>	<i>5'rpS12</i> (F)	ATTAGAAATGCAAGACAGCCAAT	<i>rpL20</i> (R)	CGTTTTTCGRGCTATA/GTATCC	Shaw <i>et al.</i> , 2005*
<i>trnC-trnD</i>	<i>trnC</i>	CCAGTTCGAATCCGGGTGTC	<i>trnD</i>	GGGATTGTAGCTCAATTGGT	Demesure <i>et al.</i> , 1995*
<i>trnT-trnF</i>	<i>trnT-F</i> (F)	CATTACAAATGCGATGCTCT	<i>trnT-F</i> (R)	ATTTGAACTGGTGACACGAG	Taberlet <i>et al.</i> , 1991*
mtDNA					
<i>nad1</i> intron2	<i>nad1</i> F <sub>1</sub>	GATCGGCCATAAATGTACTCC	<i>nad1</i> R <sub>1</sub>	CCCCATATATTCCCGGAGC	Song <i>et al.</i> , 2002
<i>nad5</i> inton 1	<i>nad5</i> -aF	GGAAATGTTGATGCTTCTTGGG	<i>nad5</i> -bR	CTGATCCAAAATCACCTACTCG	Wang <i>et al.</i> , 2000

\* Primers that have been modified in this study.

TABLE 3. Summary of sequence variation in cp- and mt-DNA regions of the genus Cedrus

Region	Ingroups length (bp)	Alignment length (bp)	Substitution sites (informative)		
			Ingroups + outgroups	Ingroups	Indels in ingroups (informative)
<i>psbB-psbH</i>	404–409	415	24 (4)	0 (0)	1 (0)
<i>rpL16</i>	799	824	82 (30)	1 (0)	0 (0)
<i>5' rps12-rpL20</i>	705	706	52 (16)	1 (0)	0 (0)
<i>trnC-trnD</i>	2253–2259	2513	368 (90)	13 (2)	3 (1)
<i>trnT-trnF</i>	1336	1417	171 (58)	3 (0)	0 (0)
cp regions combined	5497–5508	5875	697 (198)	18 (2)	4 (1)
<i>nad1</i> intron 2	530–852	622*	54 (8)	8 (0)	2 (2)
<i>nad5</i> intron 1	1252	1266	59 (28)	10 (1)	0 (0)
mt regions combined	1782–2104	1888	113 (36)	18 (1)	2 (2)

\* Some regions that could not be aligned have been removed.

by the program r8s, which summarizes the bootstrap distribution of divergence times for each node that were used to calculate standard errors.

Although fossil and geological event can both be selected as a calibration point for time estimation, the latter as the calibration point may prove to be circular. The earliest identifiable cedar fossil dates back to the Paleocene (Blokhina, 1998), and therefore a minimum date of 54.8 Myr was applied to the node connecting *Cedrus* to the outgroup in the present study.

## RESULTS

The five cpDNA fragments, *psbB-psbH*, *rpL16* intron, *5' rps12-rpL20*, *trnC-trnD* and *trnT-trnF*, are 404–409, 799, 705, 2253–2259 and 1336 bp in length, respectively (Table 3). No intraspecific sequence variation was detected. In the *trnC-trnD* region, three 5-bp gaps, one shared by *C. libani*, *C. brevifolia* and *C. atlantica*, one shared by *C. deodara* and *C. atlantica* and another one unique to *C. deodara*, were found. In addition, a 5-bp gap occurred in the *psbB-psbH* region of *C. atlantica* that included a 256-bp coding sequence. The partition-homogeneity test indicated that data sets of these cpDNA fragments were significantly congruent ( $P = 0.996$ ). The combined cpDNA data set had 5875 characters, of which 697 were variable and 198 were parsimony-informative.

The two mtDNA fragments, *nad1* intron 2 (partial sequence) and *nad5* intron 1, are 530–852 and 1252 bp in length, respectively (Table 3). No intraspecific sequence variation was detected. In the *nad1* intron 2, two deletions, relative to *C. deodara* and outgroups, were found. One of 116 bp occurred in *C. atlantica*, and the other one of 322 bp was shared by *C. libani* and *C. brevifolia*. The partition-homogeneity test indicated that data sets of the two mtDNA regions were fully congruent ( $P = 1$ ). The combined mtDNA data set was 1888 bp in length after removing some sequences that could not be aligned, including 113 variable sites, of which 36 were parsimony-informative.

Parsimony analysis of the combined cpDNA dataset generated only one most-parsimonious tree with a tree length of 743, a consistency index (CI) of 0.9960 and a retention

index (RI) of 0.9851 (tree not shown). Phylogenetic analysis of the combined mtDNA dataset also found single most-parsimonious tree (tree length = 122, CI = 0.9590, RI = 0.8780) (tree not shown). A further partition-homogeneity test between cpDNA and mtDNA data sets suggested that they were completely congruent ( $P = 1$ ). Parsimony analysis of the combined cp- and mt-DNA data set yielded a single most-parsimonious tree (tree length = 865, CI = 0.9908, RI = 0.9671) topologically the same as the cp- and mt-DNA trees, and all clades of the tree were robustly supported by the bootstrap analysis (Fig. 1). It was found that the Himalayan cedar *C. deodara* diverged first, and then the North African species *C. atlantica* separated from the common ancestor of *C. libani* and *C. brevifolia*, two species from the eastern Mediterranean area.

When an LRT compared likelihood scores under F81 (the best model given by Akaike Information Criterion), with and without the clock assumption on the ML tree obtained




FIG. 1. The single most-parsimonious tree obtained from sequence analysis of combined cp- and mt-DNA regions (length = 865, CI = 0.9908, RI = 0.9671). Numbers above the branches denote branch lengths and bootstrap values (in parentheses), respectively. Numbers below the branches are bootstrap values above 50% yielded in the cp- (left) and mt- (right) DNA trees, respectively. The present distribution of each species was indicated on the right.




FIG. 2. Cedar phylogeny and time scale of evolution determined by NPRS based on the combined chloroplast DNA data, with a calibration point of 54.8 Myr at the root node (indicated by a star) where *Cedrus* diverged from the outgroups. Standard errors for estimates of node ages are given for two nodes.

from the combined chloroplast data, the molecular clock was rejected in favour of the null hypothesis of no clock ( $\delta = 2.62614$ , d.f. = 2,  $P > 0.05$ ). Thus, a time-scaled NPRS ultrametric tree estimated under TreeEdit, using the topology with initial branch lengths estimated under maximum likelihood (Ribera *et al.*, 2004), was generated and shown in Fig. 2. On the basis of a crown group age estimate of 54.8 Myr, molecular clock estimates suggested the divergence between the North African *C. atlantica* and the eastern Mediterranean clade comprising *C. libani* and *C. brevifolia* at  $18.81 \pm 1.25$  Myr and the split between *C. libani* from Turkey, Lebanon and Syria and *C. brevifolia* from Cyprus at  $6.56 \pm 1.20$  Myr. In the PL analysis, divergence times of the above nodes were estimated to be  $23.49 \pm 3.55$  and  $7.83 \pm 2.79$  Myr, respectively.

## DISCUSSION

### *Intragenetic classification and phylogeny of Cedrus*

Based on morphological and anatomical characteristics, all taxonomists accept the Himalaya cedar *C. deodara*, but classification of the circum-Mediterranean cedars is quite controversial (Davis, 1965; Meikle, 1977; Farjón, 1990, 2001; Scaltsoyiannes, 1998). Davis (1965) treated the circum-Mediterranean cedars as a single species, *C. libani*. Tutin *et al.* (1964) and Meikle (1977) recognized the North African cedar as a separate species *C. atlantica* based on its downy young twigs and small female cones. According to a combination of morphological characters such as angle of branch, cone size, male strobili length, leaf length, number of leaves in pseudo-whorls, together with native distributions, Farjón (1990, 2001) divided the circum-Mediterranean cedars into three closely related species, i.e. *C. atlantica*, *C. brevifolia* and *C. libani*. The external morphology is often affected by biotic and environmental factors (Davis, 1965; Meikle, 1977; Farjón, 1990; Pijut, 2000), and many characters vary continuously among cedar species, which could be responsible for the difficulty of intragenetic division of *Cedrus*.

Through terpene composition analysis, Canard *et al.* (1997) found a clear differentiation between *C. libani* and

*C. atlantica*. This finding is further supported by the isozyme analysis of Scaltsoyiannes (1998), in which the dendrogram of genetic distances of 21 cedar populations comprises five distinct groups corresponding to *C. libani* ssp. *libani*, *C. libani* ssp. *stenocoma*, *C. libani* ssp. *brevifolia*, *C. deodara* and *C. atlantica*, respectively. Furthermore, fluorochrome banding patterns indicate that *C. atlantica* could be recognized as a separate species, and *C. brevifolia* as a subspecies of *C. libani*, although the three species are closely related (Bou Dagher-Kharrat *et al.*, 2001). Based on open- and controlled-pollinations, as well as analyses using nuclear and cytoplasmic markers, Fady *et al.* (2003) found that gene flow occurs naturally between *Cedrus* taxa in plantation forests, and there are no strong pre-zygotic reproductive isolating barriers among Mediterranean cedars, when sympatric. Therefore, they proposed that Mediterranean *Cedrus* should be considered as units of a single collective species comprising two regional groups, North Africa and the Middle East.

The distinctness of *C. deodara* and the close relationships among the Mediterranean cedars are further corroborated by the present cp- and mt-DNA analyses. We found that the Himalaya cedar *C. deodara* has a basal position, and *C. atlantica* is sister to the clade comprising *C. libani* and *C. brevifolia* (Fig. 1). The tree topology, together with branch length, also supports North African *C. atlantica* as a separate species, and the reciprocal monophyly of *Cedrus* species could be the result of geographical isolation, which prevents interspecific gene flow from occurring.

### *Biogeography of Cedrus*

Distribution patterns of organisms are greatly influenced by climatic and geological changes as well as their evolutionary histories. Climate tolerance is a crucial factor for plant distribution, in particular cold and aridity for trees (Pither, 2003; Svenning, 2003). From the beginning of the Tertiary, the global climate changed from extremes of expansive warmth with ice-free poles to extremes of cold with massive continental ice-sheets and polar ice caps, including several warm periods during the mid-Paleocene to early Eocene, and mid-Miocene (Zachos *et al.*, 2003). The climatic oscillation provided opportunities for biota dispersal and subdivision (Mai, 1989; LePage, 2003; Sanmartín, 2003; Spicer *et al.*, 2003; Comes, 2004; Hellwig, 2004; Vasiliev *et al.*, 2004).

Farjón (1990) proposed that the highly disjunct distribution of apparently very closely related taxa of *Cedrus* could come from contraction of a formerly more extensive range of possibly one or two species, and a possible migration route that lay across the so called Thetys Fold Belt, a series of late Tertiary mountain uplifts along the brink of the old Thetys Sea, once extending from south-east Asia to north-west Africa. However, true cedars have reliable fossil records from the Oligocene of western Kazakhstan, the Miocene of south-west Europe, the Pliocene of south-east Europe, and the early Pleistocene of the Ahaggar Massif in the central Sahara (Florin, 1963; Farjón, 1990), in addition to fossil wood from the

Paleocene of Kamchatka in Russian Far East (Blokhina, 1998, 2005). The fact that the fossils are progressively younger from north to south suggests an origin of *Cedrus* in the high latitude area of Eurasia. Its present distribution in several isolated regions could result from vicariance of southerly migrated populations during climatic oscillation in the Tertiary and further fragmentation and dispersal of these populations. This inference is robustly supported by the molecular phylogenies constructed in this study (Fig. 1), in which the Himalaya cedar is sister to the clade comprising Mediterranean cedars that was further divided into two subclades, i.e. the North African subclade and the subclade consisting of *C. libani* and *C. brevifolia*, two species from the eastern Mediterranean area.

The ancestor of Mediterranean cedars might have reached south Europe in the Miocene, which is supported by the fossil evidence mentioned above and the molecular clock estimate of  $23.49 \pm 3.55$  Myr (PL) to  $18.81 \pm 1.25$  Myr (NPRS) for the divergence between the North African *C. atlantica* and the eastern Mediterranean clade comprising *C. libani* and *C. brevifolia* (Fig. 2). The eastward dispersal of this lineage could be finally stopped by aridity in Asia Minor (Wulff, 1944; Griffin, 2002; Liu and Yin, 2002). On the other hand, biological communications between Europe and north Africa could have occurred through the Iberian connection during the Messinian (7–5 Mya) as a consequence of the desiccation of the Tethys, the Italian connection and the Arabian connection during the Miocene (Mai, 1989; Fici, 1991; Geraads, 1998; Krijgsman, 2002; Sanmartín, 2003; Hellwig, 2004). The last two connections, however, were unavailable to *Cedrus* for the presence of Saharan and Arab-Syrian deserts (Griffin, 2002; Liu and Yin, 2002; Sanmartín, 2003). It is very likely that *Cedrus* migrated into North Africa in the very late Tertiary, although *C. atlantica* was isolated from the eastern Mediterranean clade as early as the Miocene. The arrival of *Cedrus* in the Himalayas should not have been before the Miocene, when after the phased or fast uplift of the Tibetan plateau happened as suggested by recent studies, although there are a lot of debates on the time of the uplift (Dettman *et al.*, 2003; LePage, 2003; Spicer *et al.*, 2003). This inference is also corroborated by fossil pollen records from the late Miocene and Pliocene (8.10–4.07 Mya) of the central Loess Plateau in China (Ma *et al.*, 2005) and from the Pliocene of the central Himalayas (Hsü *et al.*, 1973).

It is obvious that more fossil evidence is still needed to retrieve the place of origin of *Cedrus*. In addition, the present molecular clock estimation of divergence times of true cedars is based on the fossil wood from the Paleocene, the earliest one recorded (Blokhina, 2005). The time values obtained in the present study could be younger than the real divergence times of the group, considering that *Cedrus* represents a deep or basal branch in the phylogeny of Pinaceae, a family with an origin dating back to the early Cretaceous (Wang *et al.*, 2000). To investigate in detail the differentiation process and dispersal routes of *Cedrus*, it may be necessary to conduct a phylogeographical study on all of the four cedar species in the future, with extensive population sampling.

## ACKNOWLEDGEMENTS

The authors thank the handling editor and the two anonymous reviewers for their insightful comments and suggestions on the manuscript. We also thank Prof. Zsolt Debreczy of the International Dendrological Research Institute, Wellesley, MA, USA for providing some leaf samples of *Cedrus*, especially *C. brevifolia*, a species endemic to Cyprus; Dr S. Tugrul Koruklu in University of Ankara for his kind help in collecting population samples of *C. libani*; Drs Xiao-Xin Wei, Xian-Zhao Kan, Dan Peng, Fu-Sheng Yang, Wei Wang and Rui-Qi Li for their help in the laboratory or data analysis. This study was supported by the National Natural Science Foundation of China (grant no. 30425028) and the Chinese Academy of Sciences (grant KZCX2-YW-415 and the special fund to Xiao-Quan Wang).

## LITERATURE CITED

- Ahuja MRV. 2001. Recent advances in molecular genetics of forest trees. *Euphytica* **121**: 173–195.
- Axelrod DI, Al-Shehbaz I, Raven PH. 1996. History of the modern flora of China. In: Zhang AL, Wu SG, eds. *Floristic characteristics and diversity of East Asian plants*. Beijing: Higher Education Press.
- Baum DA, Small RL, Wendel JF. 1998. Biogeography and floral evolution of baobabs (*Adansonia*, Bombacaceae) as inferred from multiple data sets. *Systematic Biology* **47**: 181–207.
- Birky CW Jr. 1995. Uniparental inheritance of mitochondrial and chloroplast genes: mechanisms and evolution. *Proceedings of the National Academy of Sciences of the USA* **92**: 11331–11338.
- Blokhina NI. 1998. Fossil wood of *Cedrus* (Pinaceae) from the Paleogene of Kamchatka. *Paleontological Journal* **32**: 532–538.
- Blokhina NI. 2005. Fossil wood from the Paleogene of Russian Far East. XVII International Botanical Congress – Abstract 6.6.2 (P99).
- Bou Dagher-Kharrat M, Grenie G, Bariteau M, Brown S, Siljak-Yakovlev S, Savouré A. 2001. Karyotype analysis reveals interspecific differentiation in the genus *Cedrus* despite genome size and base composition constancy. *Theoretical and Applied Genetics* **103**: 846–854.
- Canard D, Perru O, Tausin V, Devillard C, Bonhoure JP. 1997. Terpene composition variations in diverse provenances of *Cedrus libani* (A.) Rich. and *Cedrus atlantica* Manet. *Trees: Structure and Function* **11**: 504–510.
- Cheng WC, Fu LK. 1978. *Flora Reipublicae Popularis Sinicae*. Beijing: Science Press.
- Comes HP. 2004. The Mediterranean region – a hotspot for plant biogeographic research. *New Phytologist* **164**: 11–14.
- Cowling RM, Rundel PW, Lamont BB, Arroyo MK, Arianoutsou M. 1996. Plant diversity in Mediterranean-climate regions. *Trends in Ecology and Evolution* **11**: 362–366.
- Davis PH. 1965. *Flora of Turkey and the East Aegean islands*. Edinburgh: Edinburgh University Press.
- Demesure B, Sodji N, Petit RJ. 1995. A set of universal primers for amplification of polymorphic non-coding regions of mitochondrial and chloroplast DNA in plants. *Molecular Ecology* **4**: 129–131.
- Dettman DL, Fang XM, Garzzone CN, Li JJ. 2003. Uplift-driven climate change at 12 Ma: a long delta O-18 record from the NE margin of the Tibetan plateau. *Earth and Planetary Science Letters* **214**: 267–277.
- Efron B, Tibshirani RJ. 1993. *An introduction to the bootstrap*. Monographs on Statistics and Applied Probability No. 57. New York, NY: Chapman and Hall.
- Fady B, Lefèvre F, Reynaud M, Vendramin GG, Bou Dagher-Kharrat M, Anzidei M, *et al.* 2003. Gene flow among different taxonomic units: evidence from nuclear and cytoplasmic markers in *Cedrus* plantation forests. *Theoretical and Applied Genetics* **107**: 1132–1138.
- Farjón A. 1990. *Pinaceae: drawings and descriptions of the genera Abies, Cedrus, Pseudolarix, Keteleeria, Nothotsuga, Tsuga, Cathaya, Pseudotsuga, Larix and Picea*. Königstein: Koeltz Scientific Books.

- Farjón A. 2001. *World checklist and bibliography of conifers*, 2nd edn. London: Royal Botanic Gardens, Kew.
- Farris JS, Källersjö M, Kluge AG, Bult C. 1995. Testing significance of incongruence. *Cladistics* 10: 315–319.
- Felsenstein J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783–791.
- Felsenstein J. 1993. PHYLIP (Phylogeny Inference Package) Version 3.6a2. Distributed by the author. Department of Genome Sciences, University of Washington, Seattle. Available at <http://evolution.gs.washington.edu/phylip.html>.
- Fici S. 1991. Floristic relations between eastern Africa and the Mediterranean region with special references to northern Somalia. *Flora Mediterranea*, Vol. I, 175–185.
- Florin R. 1963. The distribution of conifer and taxad genera in time and space. *Acta Horti Bergianzo* 20: 194–256.
- Frankis MP. 1988. Generic inter-relationships in Pinaceae. *Notes from the Royal Botanic Garden, Edinburgh* 45: 527–548.
- Geraads D. 1998. Biogeography of circum-Mediterranean Miocene-Pliocene rodents; a revision using factor analysis and parsimony analysis of endemicity. *Palaeogeography, Palaeoclimatology, Palaeoecology* 137: 273–288.
- Griffin DL. 2002. Aridity and humidity: two aspects of the late Miocene climate of North Africa and the Mediterranean. *Palaeogeography, Palaeoclimatology, Palaeoecology* 182: 65–91.
- Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Hart JA. 1987. A cladistic analysis of conifers: preliminary results. *Journal of the Arnold Arboretum* 68: 269–307.
- Hellwig FHA. 2004. Centaureinae (Asteraceae) in the Mediterranean-history of ecogeographical radiation. *Plant Systematics and Evolution* 246: 137–162.
- Hewitt GM. 2004. Genetic consequences of climatic oscillations in the Quaternary. *Philosophical Transactions of the Royal Society of London. Series B. Biological Sciences* 359: 183–195.
- Hipkins VD, Krutovskii KV, Strauss SH. 1994. Organelle genomes in conifers: structure, evolution, and diversity. *Forest Genetics* 1: 179–189.
- Hsü J, Tao JR, Sun XJ. 1973. On the discovery of a *Quercus semicarpifolia* bed in mount Shisha Pangma and its significance in botany and geology. *Acta Botanica Sinica* 15: 103–119.
- Jaramillo-Correa JP, Bousquet J, Beaulieu J, Isabel N, Perron M, Bouillé M. 2003. Cross-species amplification of mitochondrial DNA sequence-tagged-site markers in conifers: the nature of polymorphism and variation within and among species in *Picea*. *Theoretical and Applied Genetics* 106: 1353–1367.
- Krijgsman W. 2002. The Mediterranean: mare nostrum of earth sciences. *Earth and Planetary Science Letters* 205: 1–12.
- Krüssmann G. 1985. *Manual of cultivated conifers*, 2nd edn. Portland, OR: Timber Press.
- Kusumi J, Tsumura Y, Yoshimaru H, Tachida H. 2002. Molecular evolution of nuclear genes in Cupressaceae, a group of conifer. *Molecular Biology and Evolution* 19: 736–747.
- Lee C, Wen J. 2004. Phylogeny of *Panax* using chloroplast *trnC-trnD* intergenic region and the utility of *trnC-trnD* in interspecific studies of plants. *Molecular Phylogenetics and Evolution* 31: 894–903.
- LePage BA. 2003. The evolution, biogeography and palaeoecology of the Pinaceae based on fossil and extant representatives. *Acta Horticulturae* 615: 29–52.
- Li LC. 1995. Studies on the karyotype and phylogeny of the Pinaceae. *Acta Phytotaxonomica Sinica* 33: 417–432.
- Li LC, Fu YX. 1995. Studies on the cytotaxonomy and the historical phyto-geography of *Cedrus* (Pinaceae). *Acta Botanica Yunnanica* 17: 41–47.
- Liu X, Yin ZY. 2002. Sensitivity of East Asian monsoon climate to the uplift of the Tibetan Plateau. *Palaeogeography, Palaeoclimatology, Palaeoecology* 183: 223–245.
- Ma Y, Wu F, Fang X, Li J, An Z, Wang W. 2005. Pollen record from red clay sequence in the central Loess Plateau between 8.10 and 2.60 Ma. *Chinese Science Bulletin* 50: 2234–2242.
- Mai DHZ. 1989. Development and regional differentiation of the European vegetation during the Tertiary. *Plant Systematics and Evolution* 162: 79–91.
- Meikle RD. 1977. *Flora of Cyprus*, Vol. I. London: Royal Botanic Gardens, Kew.
- Mogensen HL. 1996. The hows and ways of cytoplasmic inheritance in seed plants. *American Journal of Botany* 83: 383–404.
- Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GAB, Kent J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403: 853–858.
- Pijut PM. 2000. *Cedrus* – the true cedars. *Journal of Arboriculture* 26: 218–224.
- Pither J. 2003. Climate tolerance and interspecific variation in geographic range size. *Proceeding of the Royal Society of London. Series B. Biological Sciences* 270: 475–481.
- Posada D, Crandall KA. 1998. Modeltest: testing the model of DNA substitution. *Bioinformatics* 14: 817–818.
- Price RA, Olsen-Stojkovich J, Lowenstein JM. 1987. Relationships among the genera of Pinaceae: an immunological comparison. *Systematic Botany* 12: 91–97.
- Quinn CJ, Price RA, Gadek PA. 2002. Familial concepts and relationships in the conifers based on *rbcL* and *matK* sequence comparisons. *Kew Bulletin* 57: 513–531.
- Rambaut A, Charleston MA. 2002. *TreeEdit*. 1.0a10, ed. Available at <http://www.evolve.zoo.oxford.ac.uk>
- Ribera I, Nilsson AN, Vogler AP. 2004. Phylogeny and historical biogeography of Agabinae diving beetles (Coleoptera) inferred from mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* 30: 545–562.
- Rogers SO, Bendich AJ. 1988. Extraction of DNA from plant tissues. *Plant Molecular Biology (Manual)* A6: 1–10.
- Sanderson MJ. 1997. A nonparametric approach to estimating divergence times in the absence of rate constancy. *Molecular Biology and Evolution* 14: 1218–1231.
- Sanderson MJ. 2002. Estimating absolute rates of molecular evolution and divergence times: a penalized likelihood approach. *Molecular Biology and Evolution* 19: 101–109.
- Sanderson MJ. 2003. R8s: analysis of rates (“r8s”) of evolution (and other stuff), version 1.60. <http://ginger.ucdavis.edu/r8s/>.
- Sanmartín I. 2003. Dispersal vs. vicariance in the Mediterranean: historical biogeography of the Palearctic Pachydeminae (Coleoptera, Scarabaeoidea). *Journal of Biogeography* 30: 1883–1897.
- Scaltsosyiannes A. 1998. Allozyme differentiation and phylogeny of cedar species. *Silvae Genetica* 48: 61–68.
- Scarascia-Mugnozza G, Oswald H, Piussic P, Radoglou K. 2000. Forests of the Mediterranean region: gaps in knowledge and research needs. *Forest Ecology and Management* 132: 97–109.
- Schneeweiss G, Schonswetter P, Kelso S, Niklfeld H. 2004. Complex biogeographic patterns in *Androsace* (Primulaceae) and related genera: evidence from phylogenetic analyses of nuclear internal transcribed spacer and plastid *trnL-F* sequences. *Systematic Biology* 53: 856–876.
- Shaw J, Lickey EB, Beck JT, Farmer SB, Liu W, Mille J, et al. 2005. The tortoise and the hare II: relative utility of 21 noncoding chloroplast DNA sequences for phylogenetic analysis. *American Journal of Botany* 92: 142–166.
- Small RL, Ryburn JA, Cronn RC, Seelanan T, Wendel JF. 1998. The tortoise and the hare: choosing between noncoding plastome and nuclear *Adh* sequences for phylogeny reconstruction in a recently diverged plant group. *American Journal of Botany* 85: 1301–1315.
- Song BH, Wang XQ, Wang XR, Sun LJ, Hong DY, Peng PH. 2002. Maternal lineages of *Pinus densata*, a diploid hybrid. *Molecular Ecology* 11: 1057–1063.
- Spicer RA, Harris NBW, Widdowson M, Herman AB, Guo S, Valdes PJ, et al. 2003. Constant elevation of southern Tibet over the past 15 million years. *Nature* 421: 622–624.
- Svenning JC. 2003. Deterministic Plio-Pleistocene extinctions in the European cool-temperate tree flora. *Ecology Letters* 6: 646–653.
- Swofford DL. 2002. *PAUP\*: phylogenetic analysis using parsimony (\* and other methods)*, version 4.0b10. Sunderland, MA: Sinauer Associates.
- Taberlet P, Gielly L, Pautou G, Bouvet J. 1991. Universal primers for amplification of three non-coding regions of chloroplast DNA. *Plant Molecular Biology* 17: 1105–1109.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. 1997. The CLUSTAL X windows interface: flexible strategies for

- multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* **25**: 4876–4882.
- Tsumura Y, Yoshimura K, Tomaru N, Ohba K. 1995.** Molecular phylogeny of conifers using PCR-RFLP analysis of PCR-amplified specific chloroplast genes. *Theoretical and Applied Genetics* **91**: 1222–1236.
- Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, et al. 1964.** *Flora Europaea I*, Cambridge: Cambridge University Press.
- Valcárcel V, Fiz O, Vargas P. 2003.** Chloroplast and nuclear evidence for multiple origins of polyploids and diploids of *Hedera* (Araliaceae) in the Mediterranean basin. *Molecular Phylogenetics and Evolution* **27**: 1–20.
- Vasiliev I, Krijgsman W, Langereis CG, Panaiotu CE, Matenco L, Bertotti G. 2004.** Towards an astrochronological framework for the eastern Paratethys Mio-Pliocene sedimentary sequences of the FocYani basin (Romania). *Earth and Planetary Science Letters* **227**: 231–247.
- Wang XQ, Han Y, Hong DY. 1998.** A molecular systematic study of *Cathaya*, a relic genus of the Pinaceae in China. *Plant Systematics and Evolution* **213**: 165–172.
- Wang XQ, Tank DC, Sang T. 2000.** Phylogeny and divergence times in Pinaceae: evidence from three genomes. *Molecular Biology and Evolution* **17**: 773–781.
- Wang XR, Tsumura Y, Yoshimaru H, Nagasaka K, Szmidt AE. 1999.** Phylogenetic relationships of Eurasian pines (*Pinus*, Pinaceae) based on chloroplast *rbcL*, *matK*, *rpl20-rps18* spacer and *trnV* intron sequences. *American Journal of Botany* **86**: 1742–1753.
- Wei XX, Wang XQ. 2003.** Phylogenetic split of *Larix*: evidence from paternally inherited cpDNA *trnT-trnF* region. *Plant Systematics and Evolution* **239**: 67–77.
- Winkworth RC, Donoghue MJ. 2005.** *Viburnum* phylogeny based on combined molecular data: implications for taxonomy and biogeography. *American Journal of Botany* **92**: 653–666.
- Won H, Renner SS. 2003.** Horizontal gene transfer from flowering plants to *Gnetum*. *Proceedings of the National Academy of Sciences of the USA* **100**: 10824–10829.
- Wu H, Hu ZH. 1997.** Comparative anatomy of resin ducts of the Pinaceae. *Trees: Structure and Function* **11**: 135–143.
- Wulff EV. 1944.** *Historical plant geography: history of the world flora*. Waltham, MA: Chronica Botanica.
- Xiang (J) QY, Manchester SR, Homas DT, Zhang WH, Fan CZ. 2005.** Phylogeny, biogeography, and molecular dating of cornelian cherries (*Cornus*, Cornaceae): tracking Tertiary plant migration. *Evolution* **59**: 1685–1700.
- Zachos JC, Wara MW, Bohaty S, Delaney ML, Petrizzo MR, Brill A et al. 2003.** A transient rise in tropical sea surface temperature during the Paleocene-Eocene thermal maximum. *Science* **302**: 1551–1554.